

EB

# DEUTSCHES PATENTAMT

①	Sendungen des Deutschen Patentamts sind codieren an:		<b>Antrag auf Erteilung eines Patents</b>  Aktenzeichen (wird vom Deutschen Patentamt vergeben)  <b>195 42 986.9</b>	
	Herrn Patentanwalt Dr. Hans D. Boeters  Bereiteranger 15  81541 München			
②	Zeichen des Anmelders/Vertreters (max. 20 Stellen) 7865-GBF	Telefon des Anmelders/Vertreters 089/65 00 86	Datum 17. Nov. 1995	
③	Der Empfänger in Feld ① ist der <input type="checkbox"/> Anmelder <input type="checkbox"/> Zustellungsbevollmächtigte <input checked="" type="checkbox"/> Vertreter ggf. Nr. der Allgemeinen Vollmacht			
④	<b>Anmelder</b>  Gesellschaft für Biotechnologische Forschung mbH (GBF) Mascheroder Weg 1 38124 Braunschweig		<b>Vertreter</b>  Dr. Hans D. Boeters Dipl.-Ing. Robert Bauer Dr. Enno Meyer Bereiteranger 15 81541 München	
⑤	Anmeldercode-Nr.	Vertretercode-Nr.	Zustelladressecode-Nr.	
⑥	<b>Bezeichnung der Erfindung (bei Überlänge auf gesondertem Blatt - 2fach)</b>  Epothilon-Derivate und deren Verwendung			
⑦	<b>Sonstige Anträge</b> <input type="checkbox"/> Die Anmeldung ist Zusatz zur Patentanmeldung (zum Patent) → <input type="checkbox"/> Prüfungsantrag - Prüfung der Anmeldung (§ 44 Patentgesetz) <input type="checkbox"/> Recherchenantrag - Ermittlung der öffentlichen Druckschriften ohne Prüfung (§ 43 Patentgesetz) <input type="checkbox"/> Lieferung von Ablichtungen der ermittelten Druckschriften im <input type="checkbox"/> Prüfungsverfahren <input type="checkbox"/> Recherchungsverfahren <input type="checkbox"/> Aussetzung des Erteilungsbeschlusses auf _____ Monate (§ 49 Abs. 2 Patentgesetz) (Max. 15 Mon. ab Anmelde- oder Prioritätstag)		Aktenzeichen der Hauptanmeldung (des Hauptpatents)	
⑧	<b>Erklärungen</b> <input type="checkbox"/> Teilung/Ausscheidung aus der Patentanmeldung → <input type="checkbox"/> an Lizenzvergabe interessiert (unverbindlich) <input type="checkbox"/> mit vorzeitiger Offenlegung und damit freier Akteneinsicht einverstanden (§ 31 Abs. 2 Nr. 1 Patentgesetz)		Aktenzeichen der Stammanmeldung	
⑨	<input type="checkbox"/> Inländische Priorität (Datum, Aktenzeichen der Voranmeldung) <input type="checkbox"/> Ausländische Priorität (Datum, Land, Aktenz. der Voranmeldung) } bei Überlänge auf gesondertem Blatt - 2fach)			
⑩	Gebühreuzahlung in Höhe von <u>100,00</u> DM <input type="checkbox"/> Scheck ist beigelegt <input checked="" type="checkbox"/> Überweisung (nach Erhalt der Empfangsbescheinigung) <input type="checkbox"/> Gebührenmarken sind beigelegt (bitte nicht auf d. Rückseite kleben, ggf. auf gesond. Blatt)		Abbuchung von meinem/unserem Abbuchungskonto b. d. Dresdner Bank AG, München Nr.: _____	

## ⑪ Anlagen

1. \_\_\_\_\_ Vertretervollmacht
2. \_\_\_\_\_ Erfinderbenennung
3. \_\_\_\_\_ Zusammenfassung (ggf. mit Zeichnung Fig. \_\_\_\_\_)
4. \_\_\_\_\_ Seite(n) Beschreibung
5. \_\_\_\_\_ ggf. Bezugszeichenliste
6. \_\_\_\_\_ Seite(n) Patentansprüche  
Anzahl Patentansprüche \_\_\_\_\_
7. \_\_\_\_\_ Blatt Zeichnungen
8. \_\_\_\_\_ Abschnitt(en) d. Voranmeldung
9. \_\_\_\_\_

☐ Telefax vorab am \_\_\_\_\_




(Dr. Meyer)


17. November 1995/pl

Unser Zeichen: 7865


### Epothilonderivate und deren Verwendung

Die vorliegende Erfindung betrifft allgemein Epothilonderivate und deren Verwendung zur Herstellung von Arzneimitteln. Insbesondere betrifft die vorliegende Erfindung die Herstellung der Epothilonderivate der nachfolgend dargestellten allgemeinen Formeln 1 bis 7 sowie deren Verwendung zur Herstellung von therapeutischen Mitteln und Mitteln für den Pflanzenschutz.







2




3




4



5



6



7

In den vorstehenden Formeln 1 bis 7 bedeuten:

R = H, C<sub>1</sub>- bis C<sub>4</sub>-Alkyl;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> = H, C<sub>1</sub>- bis C<sub>6</sub>-Alkyl,

C<sub>1</sub>- bis C<sub>6</sub>-Acyl-Benzoyl,

C<sub>1</sub>- bis C<sub>4</sub>-Trialkylsilyl,

Benzyl,

Phenyl,

C<sub>1</sub>- bis C<sub>6</sub>-Alkoxy-,

C<sub>6</sub>-Alkyl-, Hydroxy- und Halogen-

substituiertes Benzyl bzw. Phenyl;

wobei auch zwei der Reste R<sup>1</sup> bis R<sup>5</sup> zu der Gruppierung  
-(CH<sub>2</sub>)<sub>n</sub>- mit n = 1 bis 6 zusammentreten können und es sich  
bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um  
gradkettige oder verzweigte Reste handelt.

In der Formel 1 sind X und Y entweder gleich oder verschieden  
und stehen jeweils für Halogen, OH, O-(C<sub>1</sub>- bis C<sub>6</sub>)-Acyl,  
O-(C<sub>1</sub>- bis C<sub>6</sub>)-Alkyl, O-Benzoyl.

In der Formel 3 steht X allgemein für -C(O)-, -C(S)-, -S(O)-,  
-CR<sup>1</sup>R<sup>2</sup>-, wobei R<sup>1</sup> und R<sup>2</sup> die Bedeutung haben wie oben  
angegeben, und -SiR<sub>2</sub>, wobei R die Bedeutung hat wie oben  
angegeben.

In der Formel 4 bedeutet X Sauerstoff, NOR<sup>3</sup>, N-NR<sup>4</sup>R<sup>5</sup>, und N-  
NHCONR<sup>4</sup>R<sup>5</sup>, wobei die Reste R<sup>3</sup> bis R<sup>5</sup> die oben angegebene  
Bedeutung haben.

In der Formel 5 bedeutet X Wasserstoff, C<sub>1</sub>- bis C<sub>18</sub>-Alkyl,  
C<sub>1</sub>- bis C<sub>18</sub>-Acyl, Benzyl, Benzoyl und Cinnamoyl.

Verbindungen gemäß der allgemeinen Formel 1 sind ausgehend  
von Epothilon A und B sowie von deren 3-O- und/oder 7-O-  
geschützt n Derivaten durch Öffnung des 12,13-Epoxids  
zugänglich. W rden dazu Hydrogenwasserst ffsäur n in ein m  
bevorzugt nicht wässrigen Lösungsmittel eingesetzt, wob i man

die Halogenhydrine  $X = \text{Hal}$ ,  $Y = \text{OH}$  und  $Y = \text{OH}$ ,  $Y = \text{Hal}$  erhält. Protonensäuren wie z.B. Toluolsulfonsäure und Trifluoressigsäure führen in Gegenwart von Wasser zu 12,13-Diolen, die anschließend nach Standardverfahren acyliert (z.B. mit Carbonsäureanhydriden und Pyridin oder Triethylamin/DMAP) oder alkyliert (Alkylhalogenide und Silberoxid) werden. Die 3- und 7-Hydroxygruppen können dazu vorübergehend als Formiat (Abspaltung mit  $\text{NH}_3/\text{MeOH}$ ) oder p-Methoxybenzylether (Abspaltung mit DDQ) geschützt werden.

Verbindungen gemäß der allgemeinen Formel 2 sind aus Epothilon A und B sowie deren 3-O- und/oder 7-O-geschützten Derivaten durch Reduktion, z.B. mit  $\text{NaBH}_4$  in Methanol erhältlich. Sind dabei 3-OH und/oder 7-OH reversibel geschützt, so können nach Acylierung oder Alkylierung und Entfernen der Schutzgruppen 5-O-monosubstituierte, 3,5- oder 5,7-O-disubstituierte Derivate der allgemeinen Formel 2 erhalten werden.

Umsetzungen von Epothilon A und B mit bifunktionellen elektrophilen Reagenzien, wie (Thio)Phosgen, (Thio)Carbonyldimidazol, Thionylchlorid oder Dialkylsilyldichloriden bzw. -bistriflaten ergeben Verbindungen der allgemeinen Formel 3. Als Hilfsbasen dienen dabei Pyridin, Trialkylamine, ggf. zusammen mit DMAP bzw. 2,6-Lutidin in einem nichtprotischen Lösungsmittel. Die 3,7-Acetale der allgemeinen Formel 3 entstehen durch Umacetalisierung z.B. von Dimethylacetalen in Gegenwart eines sauren Katalysators.

Verbindungen gemäß der allgemeinen Formel 4 werden aus Epothilon A und B oder ihren 3-O- und/oder 7-O-geschützten Derivaten durch Ozonolyse und reduktive Aufarbeitung, z.B. mit Dimethylsulfid, erhalten. Die C-16-Ketone können anschließend nach dem Fachmann geläufigen Standardverfahren in Oxim, Hydrazone oder Semicarbazone umgewandelt werden. Sie werden weiterhin durch Wittig-, Wittig-Horn- oder Julia- oder Petersen-Olefinierung in C-16/C-17-Olefin überführt.

Durch Reduktion der C-16-Ketogruppe, z.B. mit einem Aluminium- oder Borhydrid, sind die 16-Hydroxyderivate gemäß der allgemeinen Formel 5 erhältlich. Diese können, wenn 3-OH und 7-OH mit entsprechenden Schutzgruppen versehen sind, selektiv acyliert oder alkyliert werden. Die Freisetzung der 3-OH- und 7-OH-Gruppen erfolgt z.B. bei O-Formyl durch  $\text{NH}_3/\text{MeOH}$ , bei O-p-Methoxybenzyl durch DDQ.

Die Verbindungen der allgemeinen Formel 6 werden aus Derivaten von Epothilon A und B erhalten, bei denen die 7-OH-Gruppe durch Acyl- oder Ethergruppen geschützt ist, in dem die 3-OH-Gruppe z.B. formyliert, mesyliert oder tosyliert und anschließend durch Behandlung mit einer Base z.B. DBU eliminiert wird. Die 7-OH-Gruppe kann wie oben beschrieben freigesetzt werden.

Verbindungen der allgemeinen Formel 7 werden aus Epothilon A und B oder deren 3-OH- und 7-OH-geschützten Derivaten durch basische Hydrolyse erhalten, z.B. mit NaOH in MeOH. Die Carboxylgruppe kann mit Diazoalkanen nach Schutz der 19-OH-Gruppe durch Alkylierung in Ester umgewandelt werden.

Die Erfindung betrifft ferner Mittel für den Pflanzenschutz in Landwirtschaft, Forstwirtschaft und/oder Gartenbau, bestehend aus einer oder mehreren der vorstehend aufgeführten Epothilonderivate bzw. bestehend aus einem oder mehreren der vorstehend aufgeführten Epothilonderivate neben einem oder mehreren üblichen Träger(n) und/oder Verdünnungsmittel(n).

Schließlich betrifft die Erfindung therapeutische Mittel, bestehend aus einer oder mehreren der vorstehend aufgeführten Verbindungen oder einer oder mehreren der vorstehend aufgeführten Verbindungen neben einem oder mehreren üblichen Träger(n) und/oder Verdünnungsmittel(n). Diese Mittel können insbesondere cytotoxisch Aktivitäten entwickeln und/oder Immunsuppression bewirken, so daß sie besonders bevorzugt als Cytostatika verwendbar sind.

Die Erfindung wird im folgenden durch die Beschreibung von einigen ausgewählten Ausführungsbeispielen näher erläutert und beschrieben.

### Beispiele

#### Verbindung 1a:

20 mg (0.041 mmol) Epothilon A werden in 1 ml Aceton gelöst, mit 50 µl (0.649 mmol) Trifluoressigsäure versetzt und über Nacht bei 50 °C gerührt. Zur Aufarbeitung wird das Reaktionsgemisch mit 1 M Phosphatpuffer pH 7 versetzt und die wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natrium-sulfat getrocknet und vom Lösungsmittel befreit. Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schicht-chromatographie (Laufmittel: Dichlormethan/Aceton, 85 : 15). Ausbeute: 4 mg (19 % ) Isomer I  
4 mg (19 % ) Isomer II

#### Isomer I

R<sub>f</sub> (Dichlormethan/Aceton, 85 : 15): 0.46

IR (Film):  $\nu$  = 3440 (m, b, Sch), 2946 (s, Sch), 1734 (vs), 1686 (m), 1456 (m), 1375 (w), 1256 (s, Sch), 1190 (w, b, Sch), 1071 (m, Sch), 884 (w), 735 (w) cm<sup>-1</sup>.

MS (20/70 eV): m/e (%) = 493 (43 [M-H<sub>2</sub>O]<sup>+</sup>), 394 (47), 306 (32), 206 (30), 181 (40), 166 (72), 139 (100), 113 (19), 71 (19), 57 (24), 43 (24).

Hochauflösung: C<sub>26</sub>H<sub>39</sub>O<sub>6</sub>NS ber.: 493.2498 für [M-H<sub>2</sub>O]<sup>+</sup>  
gef.: 493.2478



Isomer II

R<sub>f</sub> (Dichlormethan/Aceton, 85 : 15): 0.22

IR (Film):  $\nu$  = 3484 (s, b, Sch), 2942 (vs, Sch), 1727 (vs), 1570 (w), 1456 (m), 1380 (m), 1265 (s), 1190 (w), 1069 (m), 975 (w),  $\text{cm}^{-1}$ .

MS (20/70 eV): m/e (%) = 493 (21 [M-H<sub>2</sub>O]<sup>+</sup>), 394 (12), 306 (46), 206 (37), 181 (63), 166 (99), 139 (100), 113 (21), 71 (23), 57 (33), 43 (28).

Hochauflösung: C<sub>26</sub>H<sub>39</sub>O<sub>6</sub>NS ber.: 493.2498 für [M-H<sub>2</sub>O]<sup>+</sup>  
gef.: 493.2475

Verbindung 1b:

55 mg (0.111 mmol) Epothilon A werden in 0.5 ml Tetrahydrofuran gelöst, mit 0.5 ml 1 N Salzsäure versetzt und 30 Minuten bei Raumtemperatur gerührt. Anschließend wird mit 1 N Phos-phatpuffer pH 7 versetzt und die wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit. Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schichtchromatographie (Laufmittel: Dichlormethan/Methanol, 90 : 10). Ausbeute: 19 mg (32 %)

R<sub>f</sub> (Dichlormethan/Methanol, 90 : 10): 0.46

IR (Film):  $\nu$  = 3441 (s, br, Sch), 2948 (s, Sch), 1725 (vs, Sch), 1462 (m), 1381 (w), 1265 (m), 1154 (w), 972 (m, br, Sch)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 210 (4.29), 248 (4.11) nm.

MS (20/70 eV): m/e (%) = 529 (13 [M<sup>+</sup>]), 494 (10), 342 (38), 306 (23), 194 (32), 164 (100), 140 (31), 113 (15), 57 (16).

Hochauflösung: C<sub>26</sub>H<sub>40</sub>O<sub>6</sub>ClNS ber.: 529.2265 für [M<sup>+</sup>],

gef.: 529.2280

**Verbindung 1c:**

25 mg (0.047 mmol) 12-Chlor-13-hydroxy-epothilon A (1b) werden in 1 ml Dichlormethan gelöst, mit 29 mg (0.235 mmol) Dimethylaminopyridin, 151  $\mu$ l (1.081 mmol) Triethylamin und 20  $\mu$ l (0.517 mmol) 98 %-iger Ameisensäure versetzt. Das Reaktionsgemisch wird mit Eis/Natriumchlorid abgekühlt. Nach Erreichen von - 15 °C werden dem Reaktionsgemisch 40  $\mu$ l (0.423 mmol) Essigsäureanhydrid zugegeben und 70 Minuten bei - 15 °C gerührt. Nach dem ein Dünnschichtchromatogramm keinen vollständigen Umsatz anzeigt, werden dem Reaktionsgemisch weitere 6 mg (0.047 mmol) Di-methylaminopyridin, 7  $\mu$ l (0.047 mmol) Triethylamin, 2  $\mu$ l 98 %-ige Ameisensäure (0.047 mmol) und 4  $\mu$ l (0.047 mmol) Essigsäureanhydrid zugesetzt und 60 Minuten gerührt.

Zur Aufarbeitung wird das Reaktionsgemisch auf Raumtemperatur erwärmt, mit 1 M Phosphatpuffer pH 7 versetzt und die wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit.

Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schichtchromatographie (Laufmittel: Dichlormethan/Aceton, 90 : 10).

Ausbeute 1c: 5 mg (18 %)

**Verbindung 1c:**

R<sub>f</sub> (Dichlormethan/Aceton, 90 : 10): 0.67

IR (Film):  $\nu$  = 3497 (w, b, Sch), 2940 (s, b, Sch), 1725 (vs), 1468 (m, b, Sch), 1379 (m), 1265 (s), 1253 (s), 1175 (vs), 972 (m, b, Sch), 737 (s)  $\text{cm}^{-1}$ .

MS (20/70 eV): m/e (%) = 613 (9 [M<sup>+</sup>]), 567 (43), 472 (63), 382 (23), 352 (21), 164 (100),

151 (33), 96 (31), 69 (17), 44  
(26).

Hochauflösung:  $C_{29}H_{40}O_6NSCl$  ber.: 613.2112 für  $[M^+]$   
gef.: 613.2131

**Verbindung 1d:**

10 mg (0.020 mmol) Epothilon B werden in 0.5 ml Tetrahydrofuran gelöst, mit 0.5 ml 1 N Salzsäure versetzt und 30 Minuten bei Raumtemperatur gerührt. Anschließend wird mit 1 M Phosphatpuffer pH 7 versetzt und die wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit. Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schicht-chromatographie (Laufmittel: Dichlormethan/Aceton, 85 : 15).

Ausbeute: 1 mg (9 %)

R<sub>f</sub> (Dichlormethan/Aceton, 85 : 15): 0.38

MS (20/70 eV): m/e (%) = 543 (3  $[M^+]$ ), 507 (14), 320 (19),  
234 (9), 194 (17), 182 (23), 164  
(100), 140 (22), 113 (14), 71  
(13).

Hochauflösung:  $C_{27}H_{42}O_6NSCl$  ber.: 543.2421 für  $[M^+]$   
gef.: 543.2405

**Verbindung 2a:**

100 mg (0.203 mmol) Epothilon A werden in 4 ml Tetrahydrofuran/1 M Phosphatpuffer pH 7 (1 : 1) gelöst und solange mit Natrium-borhydrid (150 mg = 3.965 mmol) versetzt bis das Edukt laut Dünn-schichtchromatogramm vollständig abreagiert ist. Anschließend wird mit 1 M Phosphatpuffer pH 7 verdünnt und die

wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit.

Die Reinigung des Rohproduktes erfolgt durch Kieselchromatographie (Laufmittel: Dichlormethan/Aceton, 95 : 5 - grad - nach Dichlormethan/Aceton, 85 : 15).

Ausbeute: (20 %)

R<sub>f</sub> (Dichlormethan/Aceton, 75 : 25): 0.27

IR (Film):  $\nu$  = 3413 (s, b, Sch), 2965 (vs, Sch), 1734 (vs), 1458 (m, b, Sch), 1383 (m, Sch), 1264 (s, b, Sch), 1184 (m, b, Sch), 1059 (s, Sch), 966 (s), 885 (w), 737 (m) cm<sup>-1</sup>

MS (20/70 eV): m/e (%) = 495 (6 [M<sup>+</sup>]), 477 (8), 452 (12), 394 (9), 364 (16), 306 (49), 194 (19), 178 (35), 164 (100), 140 (40), 83 (21), 55 (27).

Hochauflösung: C<sub>26</sub>H<sub>41</sub>O<sub>6</sub>NS ber.: 495.2655 für [M<sup>+</sup>]  
gef.: 495.2623

**Verbindung 3a-d (a-d sind Stereoisomere):**

100 mg (0.203 mmol) Epothilon werden in 3 ml Pyridin gelöst, mit 50 µl (0.686 mmol) Thionylchlorid versetzt und 15 Minuten bei Raumtemperatur gerührt. Anschließend wird mit 1 M Phosphatpuffer pH 7 versetzt und die wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit. Die Reinigung des Rohproduktes und Trennung der vier Stereoisomeren 3a-d erfolgt mit Hilfe der präparativen Schichtchromatographie (Laufmittel: Toluol/Methanol, 90 : 10).

**Verbindung 3a:**

Ausbeute: 4 mg (12 %)

R<sub>f</sub> (Toluol/Methanol, 90 : 10): 0.50

IR (Film):  $\nu$  = 2961 (m, b, Sch), 1742 (vs), 1701 (vs),  
1465 (m, Sch), 1389 (m, Sch), 1238 (s,  
Sch), 1210 (vs, Sch), 1011 (s, Sch), 957  
(s, b, Sch), 808 (m, Sch), 768 (s, Sch) cm<sup>-1</sup>.

UV (Methanol):  $\lambda_{\max}$  (lg  $\epsilon$ ) = 210 (4.50), 248 (4.35) nm.

MS (20/70 eV): m/e (%) = 539 (40 [M<sup>+</sup>]), 457 (22), 362 (16),  
316 (27), 222 (30), 178 (30), 164  
(100), 151 (43), 96 (38), 69 (29),  
55 (28), 43 (20).

Hochauflösung: C<sub>26</sub>H<sub>37</sub>O<sub>7</sub>NS<sub>2</sub> ber.: 539.2011 für [M<sup>+</sup>]

**Verbindung 3b:**

Ausbeute: 14 mg (13 %)

R<sub>f</sub> (Toluol/Methanol, 90 : 10): 0.44

IR (Film):  $\nu$  = 2963 (s, br, Sch), 1740 (vs), 1703 (s),  
1510 (w), 1464 (m, br, Sch), 1389 (m, Sch),  
1240 (s, br, Sch), 1142 (m), 1076 (w), 1037  
(w), 1003 (m), 945 (s, br, Sch), 806 (m,  
Sch), 775 (s), 737 (m) cm<sup>-1</sup>.

UV (Methanol):  $\lambda_{\max}$  (lg  $\epsilon$ ) = 211 (4.16), 250 (4.08) nm.

MS (20/70 eV): m/e (%) = 539 (27 [M<sup>+</sup>]), 475 (17), 322 (41),  
306 (67), 222 (16), 206 (17), 194  
(19), 178 (32), 164 (100), 151  
(33), 125 (18), 113 (15), 96  
(39), 81 (23), 64 (58), 57 (42),  
41 (19).

Hochauflösung: C<sub>26</sub>H<sub>37</sub>O<sub>7</sub>NS<sub>2</sub> ber.: 539.2011 für [M<sup>+</sup>]  
gef.: 539.1998

**Verbindung 3c:**

Ausbeute: 4 mg (4 %)

R<sub>f</sub> (Toluol/Methanol, 90 : 10): 0.38

MS (20/70 eV): m/e (%) = 539 (51 [M<sup>+</sup>]), 322 (22), 306 (53),  
222 (36), 178 (31), 164 (100), 151 (41), 96  
(25), 81 (20), 69 (26), 55 (25), 41 (25).

Hochauflösung: C<sub>26</sub>H<sub>37</sub>O<sub>7</sub>NS<sub>2</sub> ber.: 539.2011 für [M<sup>+</sup>]  
gef.: 539.2001

**Verbindung 3d:**

Ausbeute: 1 mg (1 %)

R<sub>f</sub> (Toluol/Methanol, 90 : 10): 0.33

MS (20/70 eV): m/e (%) = 539 (69 [M<sup>+</sup>]), 322 (35), 306 (51),  
222 (41), 178 (31), 164 (100), 151  
(46), 96 (31), 81 (26), 69 (34),  
55 (33), 41 (35).

Hochauflösung: C<sub>26</sub>H<sub>37</sub>O<sub>7</sub>NS<sub>2</sub> ber.: 539.2011 für [M<sup>+</sup>]  
gef.: 539.1997

**Verbindung 4a:**

10 mg (0.020 mmol) Epothilon A werden in 2 ml Dichlormethan gelöst, auf -70 °C abgekühlt und anschließend 5 Minuten mit Ozon bis zur schwachen Blaufärbung behandelt. Das resultierende Reaktionsgemisch wird anschließend mit 0.5 ml Dimethylsulfid versetzt und auf Raumtemperatur erwärmt. Zur Aufarbeitung wird das Reaktionsgemisch vom Lösungsmittel befreit und schließlich durch präparative Schichtchromatographie (Laufmittel: Dichlormethan/Aceton/Methanol, 85 : 10 : 5) gereinigt.

Ausbeute: 5 mg (64 %)

R<sub>f</sub> (Dichlormethan/Aceton/Methanol, 85 : 10 : 5): 0.61

IR (Film):  $\nu$  = 3468 (s, br, Sch), 2947 (s, br, Sch),  
1734 (vs, Sch), 1458 (w), 1380 (w), 1267  
(w), 1157 (w), 1080 (w), 982 (w) cm<sup>-1</sup>.

UV (Methanol):  $\lambda_{\max}$  (lg  $\epsilon$ ) = 202 (3.53) nm.

MS (20/70 eV): m/e (%) = 398 (2 [M<sup>+</sup>]), 380 (4), 267 (14),  
249 (17), 211 (20), 193 (26), 171  
(34), 139 (34), 111 (40), 96  
(100), 71 (48), 43 (50).

Hochauflösung: C<sub>21</sub>H<sub>34</sub>O<sub>7</sub>, ber.: 398.2305 für [M<sup>+</sup>]  
gef.: 398.2295.

#### Verbindung 6a:

10 mg (0.018 mmol) 3,7-Di-O-formyl-epothilon A werden in 1 ml Di-chlormethan gelöst, mit 27  $\mu$ l (0.180 mmol) 1,8-Diazabicyclo[5.4.0]undec-7-en (DBU) versetzt und 60 Minuten bei Raumtemperatur gerührt.

Zur Aufarbeitung wird das Reaktionsgemisch mit 1 M Natrium-dihydrogenphosphat-Puffer pH 4.5 versetzt und die wäßrige Phase viermal mit Ethylacetat extrahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit.

Nach Beseitigung des Lösungsmittel wird das resultierende Rohprodukt in 1 ml Methanol gelöst, mit 200  $\mu$ l einer ammoniakalischen Methanollösung (2 mmol NH<sub>3</sub>/ml Methanol) versetzt und über Nacht bei Raumtemperatur gerührt. Zur Aufarbeitung wird das Lösungsmittel im Vakuum entfernt.

Ausbeute: 4 mg (22 %)

R<sub>f</sub> (Dichlormethan/Aceton, 85 : 15): 0.46

IR (Film):  $\nu$  = 3445 (w, br, Sch), 2950 (vs, br, Sch),  
1717 (vs, Sch), 1644 (w), 1466 (m, Sch),  
1370 (m, Sch), 1267 (s, br, Sch), 1179  
(s, Sch), 984 (s, Sch), 860 (w), 733 (m)  
cm<sup>-1</sup>.

UV (Methanol):  $\lambda_{\max}$  (lg  $\epsilon$ ) = 210 (4.16) nm.

MS (20/70 eV): m/e (%) = 475 (28 [M<sup>+</sup>]), 380 (21), 322 (37),  
318 (40), 304 (66), 178 (31), 166  
(100), 151 (29), 140 (19), 96  
(38), 81 (20), 57 (26).

Hochauflösung: C<sub>26</sub>H<sub>37</sub>O<sub>3</sub>NS ber.: 475.2392 für [M<sup>+</sup>]  
gef.: 475.2384

**Verbindung 6b:**

50 mg (0.091 mmol) 3,7-Di-O-formyl-epothilon A (werden in 1 ml  
Dichlorethan gelöst, mit 2 ml (0.013 mol) 1,8-Diazabicyclo  
[5.4.0]undec-7-en (DBU) versetzt und 12 Stunden bei 90 °C  
gerührt.

Zur Aufarbeitung wird das Reaktionsgemisch mit 1 M Natrium-  
dihydrogenphosphat-Puffer pH 4.5 versetzt und die wäßrige  
Phase viermal mit Ethylacetat extrahiert. Die vereinigten  
organischen Phasen werden mit gesättigter Natriumchlorid-  
Lösung gewaschen, über Natriumsulfat getrocknet und vom  
Lösungsmittel befreit.

Die Reinigung des aus zwei Verbindungen bestehenden  
Rohproduktes erfolgt mittels präparativer  
Schichtchromatographie (Laufmittel: Dichlormethan/Aceton, 90 :  
10).

Ausbeute: 7 mg (15 %)

**Substanzcode**

R<sub>f</sub> (Dichlormethan/Aceton, 90 : 10): 0.62

IR (Film):  $\nu$  = 2951 (m, br, Sch), 1723 (vs), 1644 (w,  
br, Sch), 1468 (w), 1377 (w), 1271 (m,  
br, Sch), 1179 (s), 987 (m, br, Sch),  
735 (w, br, Sch) cm<sup>-1</sup>.

UV (Methanol):  $\lambda_{\max}$  (lg  $\epsilon$ ) = 210 (4.44) nm.

MS (20/70 eV): m/e (%) = 503 (68 [M<sup>+</sup>]), 408 (58), 390 (32),  
334 (25), 316 (34), 220 (21), 206  
(27), 194 (20), 181 (33), 164  
(100), 151 (34), 139 (28), 113  
(20), 96 (82), 81 (33), 67 (24),



55 (26), 43 (22).

Hochauflösung:  $C_{27}H_{37}O_6NS$  ber.: 503.2342 für  $[M^+]$   
gef.: 503.2303

**Verbindung 6c:**

5 mg (0.009 mmol) 3,7-Di-O-acetyl-epothilon werden in 1 ml Methanol gelöst, mit 150 µl einer ammoniakalischen Methanollösung (2 mmol  $NH_3$ /ml Methanol) versetzt und über Nacht bei 50 °C gerührt.

Zur Aufarbeitung wird das Lösungsmittel im Vakuum entfernt. Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schichtchromatographie (Laufmittel: Toluol/Methanol, 90 : 10).

Ausbeute: 3 mg (67 %)

$R_f$  (Dichlormethan/Aceton, 90 : 10): 0.55

IR (Film):  $\nu$  = 2934 (s, b, Sch), 1719 (vs, b, Sch),  
1641 (m), 1460 (m, Sch), 1372 (s, Sch),  
1237 (vs, b, Sch), 1179 (s, Sch), 1020  
(s), 963 (s, Sch), 737 (vs)  $cm^{-1}$ .

UV (Methanol):  $\lambda_{max}$  (lg  $\epsilon$ ) = 210 (4.33) nm.

MS (20/70 eV): m/e (%) = 517 (57  $[M^+]$ ), 422 (58), 318  
(31), 194 (20), 181 (34), 166  
(100), 151 (31), 96 (96), 81  
(32), 69 (27), 55 (29), 43 (69).

Hochauflösung:  $C_{28}H_{39}O_6NS$  ber.: 517.2498 für  $[M^+]$   
gef.: 517.2492

**Verbindung 7a:**

20 mg (0.041 mmol) Epothilon werden in 0.5 ml Methanol gelöst, mit 0.5 ml 1 N Natronlauge versetzt und 5 Minuten bei Raum-

temperatur gerührt.

Zur Aufarbeitung wird das Reaktionsgemisch mit 1 M. Phosphatpuffer pH 7 versetzt und die wäßrige Phase viermal mit Ethylacetat ex-trahiert. Die vereinigten organischen Phasen werden mit gesättigter Natriumchlorid-Lösung gewaschen, über Natriumsulfat getrocknet und vom Lösungsmittel befreit. Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schichtchromatographie (Laufmittel: Dichlormethan/Methanol, 85 : 15).

Ausbeute: 11 mg (52 %)

R<sub>f</sub> (Dichlormethan/Methanol, 85 : 15): 0.92

IR (Film):  $\nu$  = 3438 (s, br, Sch), 2971 (vs, br, Sch), 1703 (vs), 1507 (m), 1460 (s, Sch), 1383 (m, Sch), 1254 (w), 1190 (w, br, Sch), 1011 (w, br, Sch), 866 (w, br), 729 (s)  $\text{cm}^{-1}$ .

MS (20/70 eV): m/e (%) = 423 (0.1 [M<sup>+</sup>]), 323 (4), 168 (89), 140 (100), 85 (31), 57 (67).

Hochauflösung: C<sub>23</sub>H<sub>17</sub>O<sub>4</sub>NS ber.: 423.2443 für [M<sup>+</sup>]  
gef.: 423.2410

#### Verbindung 7b:

5 mg (0.009 mmol) 7-O-Acetyl-epothilon werden in 1 ml Methanol gelöst, mit 200  $\mu$ l einer ammoniakalischen Methanollösung (2 mmol NH<sub>3</sub>/ml Methanol) versetzt und zwei Tage bei 50 °C gerührt. Zur Aufarbeitung wird das Lösungsmittel im Vakuum entfernt. Die Reinigung des Rohproduktes erfolgt mit Hilfe der präparativen Schichtchromatographie (Laufmittel: Toluol/Methanol, 90 : 10).

Ausbeute: 3 mg (59 %)

R<sub>f</sub> (Dichlormethan/Methanol, 90 : 10): 0.63

IR (Film):  $\nu$  = 3441 (m, b, Sch), 2946 (s, Sch), 1732 (vs), 1600 (w), 1451 (m), 1375 (m), 1246

(s, b, Sch), 1013 (m, b, Sch)  $\text{cm}^{-1}$ .


UV (Methanol):  $\lambda_{\text{max}}$  ( $\lg \epsilon$ ) = 211 (3.75), 247 (3.59) nm.

MS (20/70 eV): m/e (%) = 567 (1 [ $\text{M}^+$ ]), 465 (4), 422 (7),  
388 (5), 194 (5), 182 (7), 168  
(65), 164 (17), 140 (100), 97  
(10), 71 (22), 43 (27).

Hochauflösung:  $\text{C}_{22}\text{H}_{43}\text{O}_8\text{NS}$  ber.: 567.2866 für [ $\text{M}^+$ ]  
gef.: 567.2849


## Patentansprüche

### 1. Epothilonderivat der Formel 1




wobei R = H, C<sub>1</sub>- bis C<sub>4</sub>-Alkyl; R<sup>1</sup>, R<sup>2</sup> = H, C<sub>1</sub>- bis C<sub>8</sub>-Alkyl, C<sub>1</sub>- bis C<sub>8</sub>-Acyl-Benzoyl, C<sub>1</sub>- bis C<sub>4</sub>-Trialkylsilyl, Benzyl, Phenyl, C<sub>1</sub>- bis C<sub>8</sub>-Alkoxy-, C<sub>8</sub>-Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl; und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt, und X und Y entweder gleich oder verschieden sind und jeweils für Halogen, OH, O-(C<sub>1</sub>- bis C<sub>8</sub>)-Acyl, O-(C<sub>1</sub>- bis C<sub>8</sub>)-Alkyl, O-Benzoyl stehen.

### 2. Epothilonderivat der Formel 2




wobei R = H, C<sub>1</sub>- bis C<sub>4</sub>-Alkyl; R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> = H, C<sub>1</sub>- bis C<sub>6</sub>-Alkyl, C<sub>1</sub>- bis C<sub>6</sub>-Acyl-Benzoyl, C<sub>1</sub>- bis C<sub>4</sub>-Trialkylsilyl, Benzyl, Phenyl, C<sub>1</sub>- bis C<sub>6</sub>-Alkoxy-, C<sub>6</sub>-Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl; und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt.

### 3. Epothilonderivat der Formel 3



wobei  $R = H$ ,  $C_1$ - bis  $C_4$ -Alkyl;  $R^1, R^2 = H$ ,  $C_1$ - bis  $C_6$ -Alkyl,  $C_1$ - bis  $C_6$ -Acyl-Benzoyl,  $C_1$ - bis  $C_4$ -Trialkylsilyl, Benzyl, Phenyl,  $C_1$ - bis  $C_6$ -Alkoxy-,  $C_6$ -Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl; und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt, und  $X$  allgemein für  $-C(O)-$ ,  $-C(S)-$ ,  $-S(O)-$ ,  $-CR^1R^2-$  und  $-SiR_2-$  steht, wobei  $R, R^1$  und  $R^2$  die Bedeutung hat wie oben angegeben.


4. Epothilonderivat der Formel 4



4

wobei  $R = H$ ,  $C_1$ - bis  $C_4$ -Alkyl;  $R^1, R^2 = H$ ,  $C_1$ - bis  $C_6$ -Alkyl,  $C_1$ - bis  $C_6$ -Acyl-Benzoyl,  $C_1$ - bis  $C_4$ -Trialkylsilyl, Benzyl, Phenyl,  $C_1$ - bis  $C_6$ -Alkoxy-,  $C_6$ -Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl; und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt,  $X$  Sauerstoff,  $NOR^3$ ,  $N-NR^4R^5$ , und  $N-NHCONR^4R^5$  bedeutet, wobei die Reste  $R^3$  bis  $R^5$  die oben angegebene Bedeutung haben.


5. Epothilonderivat der Formel 5



5


wobei R = H, C<sub>1</sub>- bis C<sub>4</sub>-Alkyl; R<sup>1</sup>, R<sup>2</sup> = H, C<sub>1</sub>- bis C<sub>6</sub>-Alkyl, C<sub>1</sub>- bis C<sub>6</sub>-Acyl-Benzoyl, C<sub>1</sub>- bis C<sub>4</sub>-Trialkylsilyl, Benzyl, Phenyl, C<sub>1</sub>- bis C<sub>6</sub>-Alkoxy-, C<sub>6</sub>-Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl; und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt, und X Wasserstoff, C<sub>1</sub>- bis C<sub>18</sub>-Alkyl, C<sub>1</sub>- bis C<sub>18</sub>-Acyl, Benzyl, Benzoyl und Cinnamoyl bedeutet.

6. Epothilonderivat der Formel 6



wobei R = H, C<sub>1</sub>- bis C<sub>4</sub>-Alkyl und R<sup>1</sup> = H, C<sub>1</sub>- bis C<sub>6</sub>-Alkyl, C<sub>1</sub>- bis C<sub>6</sub>-Acyl-Benzoyl, C<sub>1</sub>- bis C<sub>4</sub>-Trialkylsilyl, Benzyl, Phenyl, C<sub>1</sub>- bis C<sub>6</sub>-Alkoxy-, C<sub>6</sub>-Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl ist, und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt.

7. Epothilonderivat der Formel 7



wobei  $R = H$ ,  $C_1$ - bis  $C_4$ -Alkyl und  $R^1$ ,  $R^2 = H$ ,  $C_1$ - bis  $C_6$ -Alkyl,  $C_1$ - bis  $C_6$ -Acyl-Benzoyl,  $C_1$ - bis  $C_4$ -Trialkylsilyl, Benzyl, Phenyl,  $C_1$ - bis  $C_6$ -Alkoxy-,  $C_6$ -Alkyl-, Hydroxy- und Halogensubstituiertes Benzyl bzw. Phenyl; und es sich bei den in den Resten enthaltenen Alkyl- bzw. Acylgruppen um gradkettige oder verzweigte Reste handelt.

8. Mittel für den Pflanzenschutz in der Landwirtschaft und Forstwirtschaft und/oder im Gartenbau, bestehend aus einem oder mehreren der Verbindungen gemäß einem der vorangehenden Ansprüche oder einer oder mehreren dieser Verbindungen neben einem oder mehreren üblichen Träger(n) und/oder Verdünnungsmittel(n).

9. Therapeutisches Mittel, insbesondere zum Einsatz als Cytostatikum, bestehend aus einer oder mehreren der Verbindungen nach einem oder mehreren der Ansprüche 1 bis 7 oder ein r od r mehrer r der Verbindungen nach einem od r mehreren der Ansprüch 1 bis 7 neben einem od r m hreren üblichen Träger(n) und/oder Verdünnungsmittel(n).



### **Zusammenfassung**

Die vorliegende Erfindung betrifft Epothilonderivate und deren Verwendung.



BC

## GERMAN PATENT OFFICE

(1) In the address, give street, house number and, if applicable, P.O. box number	Shipments of the German Patent Office are to be directed to:		Application for granting a patent
	To: Dr. Hans D. Boeters Patent Attorney Bertelsranger 15 81541 Munich		File No. (assigned by the German Patent Office) 195 42 986.9
(2)	Reference of the Applicant/ Representative (max. 20 digits) 7865-GBF	Telephone No. of the Applicant/ Representative 089/65 00 86	Date Nov. 17, 1995
(3)	The recipient in field (1) is the <input type="checkbox"/> Applicant <input type="checkbox"/> Power of Attorney <input checked="" type="checkbox"/> Representative <span style="float: right;">If applicable, No. of general power of attorney</span>		
(4) Fill out only when different from field (1)	Applicant Gesellschaft für Biotechnologische Forschung mbH (GBF) Mascheroder Weg 1 38124 Braunschweig	Representative Dr. Hans D. Boeters Dipl. -Ing. Robert Baner Dr. Erno Meyer Bertelsranger 15 81541 Munich	
(5) If known	Application code No.	Representative code No.	Delivery address code No.
(6)	Description of the invention (if too long, submit on a separate sheet in 3 copies) Epodilone Derivatives and their use		
(7) See explanations and cost information on the reverse side	Other applications <input type="checkbox"/> The application is an addition to Patent Application (to the patent) - <input type="checkbox"/> Application for examination - examination of the application (§ 44 Patent Law) <input type="checkbox"/> Application for search - determination of published documents without examination (§ 43 of the Patent Law) Supply of copies of the determined publications in / 2 copies <input type="checkbox"/> Examination process <input type="checkbox"/> Research process <input type="checkbox"/> Deferral of the decision of granting to _____ months (§ 49 of Section 2 of the Patent Law) (Minimum 15 months from the date of application or priority)		File No. of the main Application (of the main Patent)
(8)	Explanations <input type="checkbox"/> Division/exclusion from the Patent Application - <input type="checkbox"/> Increased in granting license (without obligation) <input type="checkbox"/> With prior lease open and then agreement with free inspection of the documents (§ 31, Section 2, No. 1 of the Patent Law)		File No. of the initial Application
(9)	<input type="checkbox"/> Domestic priority (date, file number of the prior application) ) <input type="checkbox"/> Foreign priority (date, country, file number of the prior application) )		(in case of excessive length, submit on a separate sheet in two copies)
(10) Explanation and cost information on the reverse side	Payment of fees in the amount of 100.00 DM <input type="checkbox"/> Check is attached <input checked="" type="checkbox"/> Transfer (after receipt of advice of receipt) <input type="checkbox"/> Receipt stamps are attached (please do not glue on the reverse side, possibly on a separate sheet) <input type="checkbox"/> charge to my/our account at Dresdner Bank AG, Munich No.:		
(11) Attachments 1-4, each 3 copies	<input type="checkbox"/> Telefax sent previously on _____ 1. — Power of Attorney 2. — naming the inventor 3. — summary (optionally with drawings fig. ____) 4. — pages of specification 5. — optional reference number list 6. — pages of Patent Claims 7. — number of Patent Claims 8. — pages of drawings 9. — copy(ies) of prior application		

(12) Dr. Meyer




November 17, 1995/pl


Our reference: 7865

**EPOTHILONE DERIVATIVES AND THEIR USE**


The present invention concerns general epothilone derivatives and their use for the production of drugs. Especially, the present invention is concerned with the preparation of epothilone derivatives according to the general Formulas 1 to 7 given below, as well as with their use for the production of therapeutic agents and agents for plant protection.








2




3




3








5




6




7




DE 195 42 986.9



5



6



7



DE 195 42 986.9

In the above Formulas 1 to 7, the symbols have the following meanings:

$R = H, C_{1-4}\text{-alkyl};$

$R^1, R^2, R^3, R^4, R^5 = H, C_{1-6}\text{-alkyl},$

$C_{1-6}\text{-acyl-benzoyl},$

$C_{1-4}\text{-trialkylsilyl},$

$\text{benzyl},$

$\text{phenyl},$

$C_{1-6}\text{-alkoxy-},$

$C_6\text{-alkyl-}, \text{hydroxy and halogen-substituted benzyl or phenyl};$

also, two of the groups  $R^1$  to  $R^5$  may be combined to form the grouping  $-(CH_2)_n-$  with  $n = 1$  to 6 and the alkyl or acyl groups contained in the groups are either straight-chain or branched groups.

In Formula 1, X and Y are either identical or different and can stand for halogens, OH, O- $(C_{1-4})\text{-acyl}$ , O- $(C_{1-4})\text{-alkyl}$ , O-benzoyl.

In Formula 3, X generally stands for  $-C(O)-$ ,  $-C(S)-$ ,  $-S(O)-$ ,  $-CR^1R^2-$ , where  $R^1$  and  $R^2$  have the meaning given above and  $-\text{Si}R_3-$ , where R has the meaning given above.

In Formula 4, X stands for oxygen,  $\text{NOR}^3$ ,  $\text{N-NR}^4\text{R}^5$ , and  $\text{N-NHCONR}^4\text{R}^5$ , where the groups  $R^3$  to  $R^5$  have the meaning given above.

In Formula 5, X stands for hydrogen,  $C_{1-18}\text{-alkyl}$ ,  $C_{1-18}\text{-acyl}$ , benzyl, benzoyl and cinnamoyl.

Compounds according to general Formula 1 are accessible starting from epothilone A and B, as well as from their 3-O- and/or 7-O-protected derivatives by opening the 12,13-epoxide. When hydrogen halides are used for this purpose in a preferred nonaqueous solvent, the halohydrins  $X = \text{Hal}$ ,  $Y = \text{OH}$  and  $Y = \text{OH}$ ,  $Y = \text{Hal}$  are obtained. Protonic acids, for example, toluenesulfonic acid and trifluoroacetic acid, lead to 12,13-diols in the presence of water and then these are acylated subsequently according to standard methods (for example, with carboxylic acid anhydrides and pyridine or triethylamine/DMAP) or are alkylated (alkyl



DE 195 42 986.9

halides and silver oxide). For this purpose, the 3- and 7-hydroxy groups can be protected temporarily as the formate (cleaved with  $\text{NH}_3/\text{MeOH}$ ) or p-methoxybenzyl ether (cleaved with DDQ).

Compounds according to general Formula 2 are obtainable from epothilone A and B as well as from their 3-O- and/or 7-O-protected derivatives by reduction, for example, with  $\text{NaBH}_4$  in methanol. If the 3-OH and/or 7-OH groups are protected reversibly during this process, after acylation or alkylation, and removal of the protecting groups, 5-O-monosubstituted, 3,5- or 5,7-O-disubstituted derivatives according to general Formula 2 can be obtained.

Reactions of epothilone A and B with bifunctional electrophilic reagents, such as (thio)phosgene, (thio)carbonyldimidazole [sic], thionyl chloride or dialkylsilyl dichlorides or bistriflates give compounds having general Formula 3. The bases used as aids here can be pyridine, trialkylamine, optionally together with DMAP or 2,6-lutidine in an aprotic solvent. The 3,7-acetals having general Formula 3 are formed by transacetalization, for example, of dimethylacetals, in the presence of an acidic catalyst.

Compounds according to general Formula 4, obtained from epothilone A and B or from their 3-O- and/or 7-O-protected derivatives by ozonolysis and reductive processing, for example, with dimethyl sulfide. The C-16 ketones can then be converted to the oximes, hydrazones or semicarbazones according to standard methods known to the expert in the field. Furthermore, they are converted into C-16/C-17 olefins by the Wittig, Wittig-Horner, Julia or Petersen olefination method.

The 16-hydroxy derivatives according to general Formula 5 are obtainable by reduction of the C-16 keto group, for example, with aluminum hydride or borohydride. When the 3-OH and 7-OH groups are protected correspondingly, they can be acylated or alkylated selectively. The liberation of the 3-OH- and 7-OH groups is done, for example, with  $\text{NH}_3/\text{MeOH}$  in the case of O-formyl and with DDQ in the case of O-p-methoxybenzyl.

The compounds having general Formula 6 are obtained from derivatives of epothilone A and B in which the 7-OH group is protected by acyl or ether groups, in which the 3-OH group





is, for example, formylated, mesylated or tosylated and then eliminated by treatment with a base, for example, DBU. The 7-OH group can be liberated as described above.

Compounds having general Formula 7 are obtained from epothilone A and B or from their 3-OH- and 7-OH-protected derivatives by basic hydrolysis, for example, with NaOH in MeOH. After protection of the 19-OH group, the carboxyl group can be converted to the ester by alkylation with diazoalkanes.

Furthermore, the invention is concerned with means for plant protection in agriculture, forestry and/or gardening, consisting of one or several of the epothilone derivatives described above or consisting of one or several of the epothilone derivatives described above in addition to one or several of the usual carrier(s) and/or diluent(s).

Finally, the invention is concerned with therapeutic agents, consisting of one or several of the compounds listed above or of one or several of the compounds listed above in addition to one or several of the usual carrier(s) and/or diluent(s). These agents can exhibit especially cytotoxic activities and/or cause immune suppression, so that they can especially preferably be used as cytostatic agents.

The invention is explained further and described by the description of a few selected practical examples.

### Examples

Translator's note: In the infrared spectra in the examples, the general English abbreviations are used, s, m, w, vs, b, etc., except for Sch = shoulder and  $\nu$  (nu). Also, the abbreviation lg stands for log.



**Compound 1a**

20 mg (0.041 mmole) of epothilone A is dissolved in 1 mL of acetone, with 50  $\mu$ L (0.649 mmole) of trifluoroacetic acid is added and the mixture is stirred overnight at 50°C. For work-up, the reaction mixture is treated with 1 M pH 7 phosphate buffer and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed. The purification of the crude product is done with the aid of preparative layer-chromatography (solvent: dichloromethane/acetone, 85:15).

Yield: 4 mg (19%) isomer I

4 mg (19%) isomer II

**Isomer I**

R<sub>f</sub> (dichloromethane/acetone, 85:15): 0.46

IR (Film):  $\nu$  = 3440 (m, b, Sch), 2946 (s, Sch), 1734 (vs), 1686 (m), 1456 (m), 1375 (w), 1256 (s, Sch), 1190 (w, b, Sch), 1071 (m, Sch), 884 (w), 735 (w)  $\text{cm}^{-1}$ .

MS (20/70 eV): m/e (%) = 493 (43) [M-H<sub>2</sub>O]<sup>+</sup>, 394 (47), 306 (32), 206 (30), 181 (40), 166 (72), 139 (100), 113 (19), 71 (19), 57 (24), 43 (24).

High resolution: C<sub>22</sub>H<sub>30</sub>O<sub>6</sub>NS

calculated: 493.2498 for [M-H<sub>2</sub>O]<sup>+</sup>

found: 493.2478



**Isomer II****R<sub>f</sub>** (dichloromethane/acetone, 85:15): 0.22

**IR (Film):**  $\nu$  = 3484 (s, b, Sch), 2942 (vs, Sch), 1727 (vs), 1570 (w), 1456 (m), 1380 (m), 1265 (s), 1190 (w), 1069 (m), 975 (w),  $\text{cm}^{-1}$ .

**MS (20/70 eV):** m/e (%) = 493 (21,  $[\text{M}-\text{H}_2\text{O}]^+$ ), 394 (12), 306 (46), 206 (37), 181 (63), 166 (99), 139 (100), 113 (21), 71 (23), 57 (33), 43 (28).

**High resolution:**  $\text{C}_{26}\text{H}_{39}\text{O}_6\text{NS}$       calculated: 493.2498 for  $[\text{M}-\text{H}_2\text{O}]^+$   
    found: 493.2475

**Compound 1b**

Epothilone A, 55 mg (0.111 mmole), is dissolved in 0.5 mL of tetrahydrofuran, 0.5 mL of 1 N hydrochloric acid is added and the mixture is stirred for 30 minutes at room temperature. Then 1 N phosphate buffer of pH 7 is added and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed. The purification of the crude product is done with the aid of preparative layer chromatography (solvent: dichloromethane/methanol, 90:10).

Yield 1c: 19 mg (32%).



DE 195 42 986.9

R<sub>f</sub> (dichloromethane/methanol, 90:10): 0.46

IR (Film):  $\nu$  = 3441 (s, br, Sch), 2948 (s, Sch), 1725 (vs, Sch), 1462 (m), 1381 (w), 1265 (m), 1154 (w), 972 (m, br, Sch)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 210 (4.29), 248 (4.11) nm.

MS (20/70 eV): m/e (%) = 529 (13) [M<sup>+</sup>], 494 (10), 342 (38), 306 (23), 194 (32), 164 (100), 140 (31), 113 (15), 57 (16).

High resolution:  $\text{C}_{22}\text{H}_{20}\text{O}_6\text{ClNS}$  calculated: 529.2265 for [M<sup>+</sup>],  
found: 529.2280

### Compound 1c

12-Chloro-13-hydroxy-epothilone A (1b), 25 mg (0.047 mmole), is dissolved in 1 mL of dichloromethane, and then 29 mg (0.235 mmole) of dimethylaminopyridine, 151  $\mu\text{L}$  (1.081 mmole) of triethylamine and 20  $\mu\text{L}$  (0.517 mmole) of 98% formic acid are added. The reaction mixture is cooled with ice/sodium chloride. After reaching  $-15^\circ\text{C}$ , 40  $\mu\text{L}$  (0.423 mmole) of acetic anhydride is added to the reaction mixture, followed by stirring for 70 minutes at  $-15^\circ\text{C}$ . Since the thin-layer chromatogram did not show complete conversion, another 6 mg (0.047 mmole) of dimethylaminopyridine, 7  $\mu\text{L}$  (0.047 mmole) of triethylamine, 2  $\mu\text{L}$  of 98% formic acid (0.047 mmole) and 4  $\mu\text{L}$  (0.047 mmole) of acetic anhydride are added to the reaction mixture, followed by stirring for 60 minutes.

For work-up, the reaction mixture is heated to room temperature, 1 M phosphate buffer with pH 7 is added and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed.

The purification of the crude product is done with the aid of preparative layer chromatography (solvent: dichloromethane/acetone, 90:10).

Yield: 5 mg (18%).





**Compound 1c****R<sub>f</sub>** (Dichloromethane/acetone, 90-10): 0.67

**IR (Film):**  $\nu$  = 3497 (w, b, Sch), 2940 (s, b, Sch), 1725 (vs), 1468 (m, b, Sch), 1379 (m), 1265 (s), 1253 (s), 1175 (vs), 972 (m, b, Sch), 737 (s)  $\text{cm}^{-1}$ .

**MS (20/70 eV):** m/e (%) = 613 (9 [M<sup>+</sup>]), 567 (43), 472 (63), 382 (23), 352 (21), 164 (100), 151 (33), 96 (31), 69 (17), 44 (26).

**High resolution:**  $\text{C}_{29}\text{H}_{40}\text{O}_2\text{NSCl}$  calculation: 613.2112 for [M<sup>+</sup>]  
found: 613.2131

**Compound 1d**

10 mg (0.020 mmole) of epothilone B is dissolved in 0.5 mL of tetrahydrofuran, then 0.5 mL of 1 N hydrochloric acid is added and the mixture stirred for 30 minutes at room temperature. Then 1 M phosphate buffer at pH 7 is added and the aqueous phase is extracted 4 times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed.

The purification of the crude product is done with the aid of preparative layer chromatography (solvent: dichloromethane/acetone, 85:15).

Yield: 1 mg (9%)



**DE 195 42 986.9**

R. (Dichloromethane/acetone, 85:15): 0.38

MS (20/70 eV): m/e (%) = 543 (3, [M<sup>+</sup>]), 507 (14), 320 (19), 234 (9), 194 (17), 182 (23), 164 (100), 140 (22), 113 (14), 71 (13).

**High resolution:** C<sub>27</sub>H<sub>30</sub>O<sub>6</sub>NSCl      calculated: 543.2421 for [M]<sup>+</sup>  
found: 543.2405

### Compound 2a

Epothilone A, 100 mg (0.203 mmole), is dissolved in 4 mL of tetrahydrofuran/1 M phosphate buffer, pH 7 (1:1) and sodium borohydride (150 mg = 3.965 mmole) is added until the thin-layer chromatogram shows that the starting material reacted completely. Then the mixture is diluted with 1 M phosphate buffer, pH 7 and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed.

The purification of the crude product is done by silica gel chromatography (solvent: dichloromethane/acetone, 95:5 - in 5 steps to dichloromethane/acetone, 85:15).

**Yield: (20%)**



R. (Dichloromethane/acetone, 75:25): 0.27

IR (Film):  $\nu$  = 3413 (s, b, Sch), 2965 (vs, Sch), 1734 (vs), 1458 (m, b, Sch), 1383 (m, Sch), 1264 (s, b, Sch), 1184 (m, b, Sch), 1059 (s, Sch), 966 (s), 885 (w), 737 (m)  $\text{cm}^{-1}$

MS (20/70 eV): m/e (%) = 495 (6 [M<sup>+</sup>]), 477 (8), 452 (12), 394 (9), 364 (16), 306 (49), 194 (19), 178 (35), 164 (100), 140 (40), 83 (21), 55 (27).

High resolution:  $\text{C}_{26}\text{H}_{41}\text{O}_6\text{NS}$  calculated: 495.2655 for [M<sup>+</sup>]  
found: 495.2623

**Compound 3a-d (a-d are stereoisomers)**

Epothilone, 100 mg (0.203 mmole) is dissolved in 3 mL of pyridine, with 50  $\mu\text{L}$  (0.686 mmole) of thionyl chloride added and the mixture is stirred for 15 minutes at room temperature. Then, 1 M phosphate buffer, pH 7, is added and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed. The purification of the crude product and separation of the four stereoisomers 3a-d is done with the aid of preparative layer chromatography (solvent: toluene/methanol, 90:10).

**Compound 3a**

Yield: 4 mg (12%)



R<sub>f</sub> (toluene/methanol, 90:10): 0.50

IR (Film):  $\nu$  = 2961 (m, br, Sch), 1742 (vs), 1701 (vs),  
1465 (m, Sch), 1389 (m, Sch), 1238 (s,  
Sch), 1210 (vs, Sch), 1011 (s, Sch), 957  
(s, br, Sch), 808 (m, Sch), 768 (s, Sch)  $\text{cm}^{-1}$

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 210 (4.50), 248 (4.35) nm.

MS (20/70 eV): m/e (%) = 539 (40 [M<sup>+</sup>]), 457 (22), 362 (16),  
316 (27), 222 (30), 178 (30), 164  
(100), 151 (43), 96 (38), 69 (29),  
55 (28), 43 (20).

High resolution:  $\text{C}_{26}\text{H}_{37}\text{O}_7\text{NS}_2$  calculated: 539.2011 for [M<sup>+</sup>]

**Compound 3b**

Yield: 14 mg (13%)

R<sub>f</sub> (toluene/methanol, 90:10): 0.44

IR (Film):  $\nu$  = 2963 (s, br, Sch), 1740 (vs), 1703 (s),  
1510 (w), 1464 (m, br, Sch), 1389 (m, Sch),  
1240 (s, br, Sch), 1142 (m), 1076 (w), 1037  
(w), 1001 (m), 945 (s, br, Sch), 806 (m,  
Sch), 775 (s), 737 (m)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 211 (4.16), 250 (4.08) nm.

MS (20/70 eV): m/e (%) = 539 (27 [M<sup>+</sup>]), 475 (17), 322 (41),  
306 (67), 222 (16), 206 (17), 194  
(19), 178 (32), 164 (100), 151  
(33), 125 (18), 113 (15), 96  
(39), 81 (23), 64 (58), 57 (42),  
43 (18).





**High resolution:** C<sub>28</sub>H<sub>37</sub>O<sub>7</sub>NS<sub>2</sub>      calculated: 539.2011 for [M]<sup>+</sup>  
found: 539.1998

### Compound 3c

**Yield: 4 mg (4%)**

**R. (toluene/methanol, 90:10): 0.38**

MS (20/70 eV): m/e (I) = 539 (22), 322 (22), 306 (53), 222 (36), 178 (33), 164 (100), 151 (41), 96 (25), 81 (20), 69 (26), 55 (25), 41 (25):

**High resolution:** C<sub>26</sub>H<sub>27</sub>O<sub>7</sub>NS<sub>2</sub>      calculated: \$39.2011 for [M<sup>+</sup>]  
found: \$39.2001

### Compound 3d

**Yield:** 1 mg (1%)

**R<sub>f</sub> (toluene/methanol, 90:10): 0.33**

MS (20/70 CV): = M/a (%) = 539 (45) (M1), 322 (35), 306 (51),  
222 (41), 178 (31), 164 (100), 151  
(46), 96 (31), 81 (26), 69 (34),  
55 (33), 41 (35).

**High resolution:** C<sub>26</sub>H<sub>37</sub>O<sub>7</sub>NS<sub>2</sub>      calculated: \$39.2011 for [M<sup>+</sup>]  
found: \$39.1997



**Compound 4a**

Epothilone A, 10 mg (0.020 mmole), is dissolved in 2 mL of dichloromethane, cooled to  $-70^{\circ}\text{C}$  and then treated with ozone for 5 minutes until a weak blue coloration develops. The resulting reaction mixture is then treated with 0.5 mL of dimethyl sulfide and heated to room temperature. In the work-up, the solvent is removed from the reaction mixture and finally the product is purified with preparative layer chromatography (solvent: dichloromethane/acetone/methanol, 85:10:5).

Yield: 5 mg (64%)

R<sub>f</sub> (Dichloromethane/acetone/methanol, 85:10:5): 0.61

IR (Film):  $\nu = 3468$  (s, br, Sch),  $2947$  (s, br, Sch),  $1734$  (vs, Sch),  $1458$  (w),  $1380$  (w),  $1267$  (w),  $1157$  (w),  $1080$  (w),  $982$  (w)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg e) =  $202$  (3.53) nm.

MS (20/70 eV):  $m/e$  (%) =  $398$  (2 [M<sup>+</sup>]),  $380$  (4),  $267$  (14),  $249$  (17),  $211$  (20),  $193$  (26),  $171$  (34),  $139$  (34),  $111$  (40),  $96$  (100),  $71$  (48),  $43$  (50).

High resolution:  $\text{C}_{21}\text{H}_{34}\text{O}_7$  calculated:  $398.2305$  for [M<sup>+</sup>]  
found:  $398.2295$

**Compound 6a**

3,7-Di-O-formyl-epothilone A, 10 mg (0.018 mmole), is dissolved in 1 mL of dichloromethane,  $27\ \mu\text{L}$  (0.180 mmole) of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) is added and the mixture stirred at room temperature for 60 minutes.

For work-up, the reaction mixture is treated with 1 M sodium dihydrogen phosphate buffer, pH 4.5, and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed.



After elimination of the solvent, the resulting crude product is dissolved in 1 mL of methanol, treated with 200  $\mu$ L of ammoniacal methanol solution (2 mmole of  $\text{NH}_3$ /mL of methanol) and stirred overnight at room temperature. For work-up, the solvent is removed in vacuum.

Yield: 4 mg (22%)

R<sub>f</sub> (Dichloromethane/acetone, 85:15): 0.46

IR (Film):  $\nu$  = 3445 (w, br, Sch), 2950 (vs, br, Sch),  
1717 (vs, Sch), 1644 (w), 1466 (m, Sch),  
1370 (m, Sch), 1267 (s, br, Sch), 1179  
(s, Sch), 984 (s, Sch), 860 (w), 733 (m)  
 $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 210 (4.16) nm.

MS (20/70 eV): m/e (%) = 475 (38 [M<sup>+</sup>]), 380 (21), 322 (37),  
318 (40), 304 (66), 178 (31), 166  
(100), 151 (29), 140 (19), 96  
(38), 81 (20), 57 (26).

High resolution:  $\text{C}_{26}\text{H}_{37}\text{O}_5\text{NS}$  calculated: 475.2392 for [M<sup>+</sup>]  
found: 475.2384

#### Compound 6b

3,7-Di-O-formyl-epothilone A, 50 mg (0.091 mmole), is dissolved in 1 mL of dichloroethane, 2 mL (0.013 mole) of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) is added and the mixture stirred for 12 hours at 90°C.

For work-up, the reaction mixture is treated with 1 M sodium dihydrogen phosphate buffer, pH 4.5, and the aqueous phase extracted four times with ethyl acetate. The combined

<sup>1</sup> [Note: Compound 5 was not mentioned in the original. It jumps from 4 to 6.]



For work-up, the reaction mixture is treated with 1 M sodium dihydrogen phosphate buffer, pH 4.5, and the aqueous phase extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent removed.

The purification of the crude product, which consists of two compounds, is done with the aid of preparative layer chromatography (solvent: dichloromethane/acetone, 90:10).

Yield: 7 mg (15%)

Substance code

R. (Dichloromethane/acetone, 90:10): 0.62

IR (Film):  $\nu$  = 2951 (m, br, Sch), 1723 (vs), 1644 (w, br, Sch), 1468 (w), 1377 (w), 1271 (m, br, Sch), 1179 (s), 987 (m, br, Sch), 735 (w, br, Sch)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 210 (4.44) nm.

MS (20/70 eV): m/e (%) = 503 (68 [M<sup>+</sup>]), 408 (58), 390 (32), 334 (25), 316 (34), 220 (21), 206 (27), 194 (20), 181 (33), 164 (100), 151 (34), 139 (28), 113 (20), 95 (82), 81 (33), 67 (24), 55 (24), 43 (22).

High resolution:  $\text{C}_{27}\text{H}_{37}\text{O}_6\text{NS}$  calculated: 503.2342 for [M<sup>+</sup>]  
found: 503.2303

#### Compound 6c

3,7-Di-O-acetyl-epothilone, 5 mg (0.009 mmole), is dissolved in 1 mL of methanol, 150  $\mu\text{L}$  of an ammoniacal methanol solution (2 mmole of  $\text{NH}_3/\text{mL}$  of methanol) is added and the mixture stirred overnight at 50°C.





For work-up, the solvent is removed in vacuum and the crude product is purified with the aid of preparative layer chromatography (solvent: toluene/methanol, 90:10).

Yield: 3 mg (67%)

R<sub>f</sub> (Dichloromethane/acetone, 90:10): 0.55

IR (Film):  $\nu$  = 2934 (s, b, Sch), 1719 (vs, b, Sch), 1641 (m), 1460 (m, Sch), 1372 (s, Sch), 1237 (vs, b, Sch), 1179 (s, Sch), 1020 (s), 963 (s, Sch), 737 (vs)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 210 (4.33) nm.

MS (20/70 eV): m/e (%) = 517 (57 [M<sup>+</sup>]), 422 (58), 318 (31), 194 (20), 181 (34), 166 (100), 151 (31), 96 (96), 81 (32), 69 (27), 55 (29), 43 (69).

High resolution:  $\text{C}_{28}\text{H}_{39}\text{O}_6\text{NS}$  calculated: 517.2498 for [M<sup>+</sup>]  
found: 517.2492

#### Compound 7a

Epothilone, 20 mg (0.041 mmole), is dissolved in 0.5 mL of methanol, 0.5 mL of 1 N sodium hydroxide is added and the mixture stirred for 5 minutes at room temperature.

For work-up, the reaction mixture is treated with 1 M phosphate buffer, pH 7, and the aqueous phase is extracted four times with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulfate and the solvent is removed. The purification of the crude product is done with the aid of preparative layer chromatography. (Solvent: dichloromethane/methanol, 85:15).

Yield: 11 mg (52%)



R<sub>f</sub> (Dichloromethane/methanol, 85:15): 0.92

IR (Film):  $\nu$  = 3438 (s, br, Sch), 2971 (vs, br, Sch), 1703 (vs), 1507 (m), 1460 (s, Sch), 1383 (m, Sch), 1254 (w), 1190 (w, br, Sch), 1011 (w, br, Sch), 866 (w, br), 729 (s)  $\text{cm}^{-1}$ .

MS (20/70 eV):  $m/e$  (%) = 423 (0.1 [M<sup>+</sup>]), 323 (4), 168 (89), 140 (100), 85 (31), 57 (67).

High resolution:  $\text{C}_{22}\text{H}_{27}\text{O}_6\text{NS}$  calculated: 423.2443 for [M<sup>+</sup>]  
found: 423.2410

### Compound 7b

5 mg (0.009 mmole) of 7-O-acetyl-epothilone is dissolved in 1 mL of methanol, 200  $\mu\text{L}$  of an ammoniacal methanol solution (2 mmole of  $\text{NH}_3/\text{mL}$  of methanol) is added and the mixture is stirred for 2 days at 50°C. For work-up, the solvent is removed in vacuum. The purification of the crude product is done with the aid of preparative layer chromatography (solvent: toluene/methanol, 90:10).

Yield: 3 mg (59%)

R<sub>f</sub> (Dichloromethane/methanol, 90:10): 0.63

IR (Film):  $\nu$  = 3441 (m, b, Sch), 2946 (s, Sch), 1732 (vs), 1600 (w), 1451 (m), 1375 (m), 1246 (s, b, Sch), 1013 (m, b, Sch)  $\text{cm}^{-1}$ .

UV (Methanol):  $\lambda_{\text{max}}$  (lg  $\epsilon$ ) = 211 (3.75), 247 (3.59) nm.


MS (20/70 eV):  $m/e$  (%) = 567 (1 [M<sup>+</sup>]), 465 (4), 422 (7), 388 (5), 194 (5), 182 (7), 168 (65), 164 (17), 140 (100), 97 (10), 71 (22), 43 (27).

High resolution:  $\text{C}_{22}\text{H}_{26}\text{O}_6\text{NS}$  calculated: 567.2866 for [M<sup>+</sup>]  
found: 567.2849



## Patent Claims


1. Epothilone derivative having Formula 1



where  $R = H, C_{1-4}$ -alkyl;  $R^1, R^2 = H, C_{1-6}$ -alkyl,  $C_{1-6}$ -acyl, benzoyl,  $C_{1-4}$ -trialkylsilyl, benzyl, phenyl,  $C_{1-6}$ -alkoxy,  $C_6$ -alkyl-, hydroxy- and halogen-substituted benzyl or phenyl; and the alkyl and acyl groups in these groups are straight-chain or branched groups and X and Y are either the same or different and stand for halogen, OH,  $O-(C_{1-6})$ -acyl,  $O-(C_{1-6}$ -alkyl or O-benzoyl).




## 2. Epothilone derivative having formula 2



where R = H, C<sub>1-4</sub>-alkyl; R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> = H, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-acyl, benzoyl, C<sub>1-4</sub>-trialkylsilyl, benzyl, phenyl, C<sub>1-6</sub>-alkoxy-, C<sub>6</sub>-alkyl-, hydroxy- and halogen-substituted benzyl or phenyl; the alkyl and acyl groups contained in these groups are straight-chain or branched groups.

## 3. Epothilone derivative according to formula 3




where R = H, C<sub>1-4</sub>-alkyl; R<sup>1</sup>, R<sup>2</sup> = H, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-acyl, benzoyl, C<sub>1-4</sub>-trialkylsilyl, benzyl, phenyl, C<sub>1-6</sub>-alkoxy-, C<sub>6</sub>-alkyl-, hydroxy-, and halogen-substituted benzyl and phenyl;






the alkyl and acyl groups contained in these groups are straight-chain or branched groups and X stands generally for  $-C(O)-$ ,  $-C(S)-$ ,  $-S(O)-$ ,  $-CR^1R^2-$  and  $-SiR_2-$ , where R,  $R^1$  and  $R^2$  have the meaning given above.

4. Epothilone derivative according to formula 4



where R = H,  $C_{1-6}$ -alkyl;  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$  = H,  $C_{1-6}$ -alkyl,  $C_{1-6}$ -acyl, benzoyl,  $C_{1-6}$ -trialkylsilyl, benzyl, phenyl,  $C_{1-6}$ -alkoxy-,  $C_6$ -alkyl-, hydroxy- and halogen-substituted benzyl or phenyl; the alkyl and acyl groups contained in these groups are straight-chain or branched groups; X stands for oxygen,  $NOR^3$ ,  $N-NR^4R^5$ , and  $N-NHCONR^4R^5$ , where the groups  $R^3$  to  $R^5$  have the meaning given above.


5. Epothilone derivative having formula 5






where  $R = H, C_{1-4}$ -alkyl;  $R^1, R^2 = H, C_{1-6}$ -alkyl,  $C_{1-6}$ -acyl, benzoyl,  $C_{1-4}$ -trialkylsilyl, benzyl, phenyl,  $C_{1-6}$ -alkoxy-,  $C_6$ -alkyl-, hydroxy- and halogen-substituted benzyl or phenyl; the alkyl and acyl groups contained in these groups are straight-chain or branched groups and  $X$  stands for hydrogen,  $C_{1-18}$ -alkyl,  $C_{1-18}$ -acyl, benzyl, benzoyl and cinnamoyl.

6. Epothilone derivative according to formula 6



in which  $R = H, C_{1-4}$ -alkyl and  $R^1 = H, C_{1-6}$ -alkyl,  $C_{1-6}$ -acyl, benzoyl,  $C_{1-4}$ -trialkylsilyl, benzyl, phenyl,  $C_{1-6}$ -alkoxy-,  $C_6$ -alkyl-, hydroxy- and halogen-substituted benzyl or phenyl; the alkyl and acyl groups contained in these groups are straight-chain or branched groups.

7. Epothilone derivative according to formula 7





in which R = H, C<sub>1-4</sub>-alkyl; and R<sup>1</sup>, R<sup>2</sup>, = H, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-acyl, benzoyl, C<sub>1-4</sub>-trialkylsilyl, benzyl, phenyl, C<sub>1-6</sub>-alkoxy-, C<sub>6</sub>-alkyl-, hydroxy- and halogen-substituted benzyl or phenyl; the alkyl and acyl groups contained in these groups are straight-chain or branched groups.

8. Means for plant protection in agriculture and forestry and/or in gardening, consisting of one or several of the compounds according to one of the previous Claims, or according to one or several of these compounds together with one or several usual carrier(s) and/or diluent(s).

9. Therapeutic agent, especially for use as cytostatic agent, consisting of one or several of the compounds according to one or several of Claims 1 to 7, or one or several compounds according to one or several of Claims 1 to 7 together with one or several of the usual carrier(s) and/or diluent(s).

#### Summary

The present invention is concerned with epothilone derivatives and their application.

